

| | INDICATOR | DESCRIPTION |
|---|---|---|
| Core RMNCAH-N Impact Indicators | Maternal Mortality Ratio | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_MMR.pdf |
| | Under 5 Mortality Ratio | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_U5MR.pdf |
| | Neonatal Mortality Ratio | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_NMR.pdf |
| | Stillbirths | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_stillbirth.pdf |
| | Adolescent Birth Rate (15-19yrs) | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_ABR.pdf |
| | Percent of births <24 months after the preceding birth | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_births_less_than_24_months_after_preceding_birth.pdf |
| | Prevalence of stunting among children under 5 years of age | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_stunting.pdf |
| | Moderate to severe wasting among children under 5 years of age | https://data.gffportal.org/sites/live/files/2024-02/GFF_reference_sheet_wasting.pdf |
| | Core Health Financing Indicators | Domestic General Government Health Expenditure (DGGHE) as share of General Government Expenditure (GGE) (%) |
| Domestic General Government Health Expenditure (DGGHE) per capita (US\$) | | Government health expenditure per inhabitant (numerator: Total government health expenditures; denominator: Total Population) |
| Out-of-pocket spending on health, per capita (US\$) | | Sum of household out-of-pocket payments on health, per capita (US\$) Share of government budget allocated to health (%) |
| Share of the government budget allocated to health | | (numerator: total government budget allocated to health; denominator: total government budget) |
| Health budget execution (%) | | Share of government health spending to total government health budget (numerator: Total government spending on health; denominator: total government budget allocated to health). |
| Share of health expenditure going to frontline providers (%) | | Share of domestic government health spending to frontline providers (country-specific definition) to total government health spending (numerator: total domestic government expenditure on frontline providers (country definition); denominator: total government expenditure on health). This indicator is not comparable across countries. |
| | Demand for FP satisfied by Modern Methods | Percentage of demand for family planning satisfied by modern methods is calculated as the number of all women using modern methods of family planning divided by the number of all women with demand for family planning (either with unmet need or currently using any family planning) |
| | Antenatal Care: Received Iron supplementation for 90+ days | Percentage of women with a birth in the five years preceding the survey who took iron tablets or syrup for 90+ days |
| | Antenatal Coverage 4+ visits | Percentage of women who had a live birth in the five (or three) years preceding the survey who had 4+ antenatal care visits |
| | Care Seeking for Pneumonia | Percentage of children under age 5 who had symptoms of ARI in the last two weeks, for whom advice or treatment was sought from a health facility or provider. |

Indicator_Definitions

RMNCAH-N Coverage Indicators

| | |
|--|---|
| DPT3 coverage | Percentage of children 12-23 months who had received DPT3 vaccination as reported by either mothers recall or the vaccination card |
| Institutional Delivery | Percentage of live births in the five (or three) years preceding the survey delivered at a health facility |
| Exclusive Breast Feeding | Percentage of children between 0 to 5 months of age living with the mother who are exclusively breastfed |
| Minimum Adequate Diet | Number of children 6-23 months that satisfy the Minimum dietary diversity and Minimum Meal Frequency requirement for breast fed and non-breast fed children. |
| Oral rehydration salts treatment of diarrhea | Percentage of children born in the five (or three) years preceding the survey with diarrhea in the two weeks preceding the survey who received oral rehydration solution (ORS), that is either fluid from an ORS packet or a pre-packaged ORS fluid. |
| Postnatal care for mothers | Percentage of women with a live birth in the 2 years preceding the survey who received a postnatal check during the first 2 days after giving birth |
| Skilled Attendance at Birth | Percentage of live births in the five (or three) years preceding the survey assisted by a skilled provider. Skilled provider includes doctor, nurse, midwife and auxiliary nurse or midwife. |
| Vitamin A supplementation, full Coverage | Percentage of children age 6-59 months who received at least one high dose Vitamin A supplement in the last 6 months |
| Adjusted net enrollment rate; lower secondary; both sexes (%) | Adjusted net enrollment is the number of pupils of the school-age group for lower secondary education, enrolled in lower secondary education, expressed as a percentage of the total population in that age group. |
| Adjusted net enrollment rate; lower secondary; female (%) | Adjusted net enrollment is the number of female pupile of the school-age group for lower secondary education, enrolled in lower secondary education, expressed as a percentage of the total population of females in that age group. |
| Lower secondary completion rate; both sexes (%) | Lower secondary education completion rate is measured as the gross intake ratio to the last grade of lower secondary education (general and pre-vocational). It is calculated as the number of new entrants in the last grade of lower secondary education, regardless of age, divided by the population at the entrance age for the last grade of lower secondary education. |
| Lower secondary completion rate; female (%) | Lower secondary education completion rate is measured as the gross intake ratio to the last grade of lower secondary education (general and pre-vocational). It is calculated as the number of new female entrants in the last grade of lower secondary education, regardless of age, divided by the female population at the entrance age for the last grade of lower secondary education. |
| SP/Fansidar 3+ doses: | Percentage of women who, during the pregnancy that resulted in the last live birth, received one or more doses of SP/Fansidar, received two or more doses of SP/Fansidar, and who received three or more doses of SP/Fansidar |